



Antibiotic and nematocidal metabolites from two lichen species collected on the island of Lampedusa (Sicily)

Jesús García Zorrilla^{1,2,*}, Trifone D'Addabbo³, Emanuela Roscetto⁴, Chiara Varriale⁴, Maria Rosaria Catania⁴, Maria Chiara Zonno⁵, Claudio Altomare⁵, Giuseppe Surico⁶, Pier Luigi Nimis⁷, Antonio Evidente¹

¹ Department of Chemical Sciences, University of Naples "Federico II", Complesso Universitario Monte Sant'Angelo, Via Cintia 4, 80126 Napoli, Italy; antonio.evidente@unina.it (A.E.)

² Allelopathy Group, Department of Organic Chemistry, Institute of Biomolecules (INBIO), School of Science, University of Cadiz, C/Republic Saharaui, s/n, 11510, Puerto Real, Cadiz, Spain

³ Institute for Sustainable Plant Protection-CNR, Unit of Bari, via G. Amendola 122/d, Bari 70126, Italy. trifone.daddabbo@ipsp.cnr.it (TD)

⁴ Department of Molecular Medicine and Medical Biotechnologies, University of Naples Federico II, Via Pansini 5, 80131 Napoli, Italy; emanuela.roscetto@unina.it (E.R.); mariarosaria.catania@unina.it (M.R.C.); chiaravariale98@gmail.com (C.V.)

⁵ Institute of Sciences of Food Production Sciences, National Research Council, Via Amendola 122/O, 70125 Bari, Italy; mariachiara.zonno@ispa.cnr.it (M.C.Z.); [claudio.altomare@ispa.cnr.it\(C.A.\)](mailto:claudio.altomare@ispa.cnr.it(C.A.))

⁶ Department of Agriculture, Food, Environment, and Forestry (DAGRI), Section of Agricultural Microbiology, Plant Pathology and Entomology, University of Florence, Italy Italy giuseppe.surico@unifi.it (G.S.)

⁷ Department of Life Sciences, University of Trieste, Via L. Giorgieri 10, 34127 Trieste, Italy; nimis@units.it (P.L.N.)

* Correspondence: e-mail: jesus.zorrilla@uca.es (J.G.Z.)



Index

Figure S1. ^1H NMR spectrum (400 MHz) of (+)-usnic acid (1) in CDCl_3	3
Figure S2. ESI MS spectrum of (+)-usnic acid (1) recorded in positive mode	3
Figure S3. ^1H NMR spectrum (500 MHz) of methyl orsellinate (2) in CDCl_3	4
Figure S4. ESI MS spectrum of methyl orsellinate (2) recorded in positive mode	4
Figure S5. ^1H NMR spectrum (500 MHz) of orcinol (3) in CDCl_3	5
Figure S6. ESI MS spectrum of orcinol (3) recorded in positive mode	5
Figure S7. ^1H NMR spectrum (400 MHz) of (+)-montagnetol (4) in CD_3OD	6
Figure S8. ESI MS spectrum of (+)-montagnetol (4) recorded in positive mode.....	6
Figure S9. ^1H NMR spectrum (400 MHz) of 4-cloroorcinol (5) in CDCl_3	7
Figure S10. ESI MS spectrum of 4-cloroorcinol (5) recorded in positive mode.....	7




Figure S1. ^1H NMR spectrum (400 MHz) of (+)-usnic acid (**1**) in CDCl_3




Figure S2. ESI MS spectrum of (+)-usnic acid (**1**) recorded in positive mode




Figure S3. ¹H NMR spectrum (500 MHz) of methyl orsellinate (**2**) in CDCl₃




Figure S4. ESI MS spectrum of methyl orsellinate (**2**) recorded in positive mode




Figure S5. ¹H NMR spectrum (500 MHz) of orcinol (**3**) in CDCl_3




Figure S6. ESI MS spectrum of orcinol (**3**) recorded in positive mode




Figure S7. ^1H NMR spectrum (400 MHz) of (+)-montagnetol (**4**) in CD_3OD




Figure S8. ESI MS spectrum of (+)-montagnetol (**4**) recorded in positive mode




Figure S9. ^1H NMR spectrum (400 MHz) of 4-chloroorcinol (**5**) in CDCl_3




Figure S10. ESI MS spectrum of 4-chloroorcinol (**5**) recorded in positive mode