

A preliminary checklist of the species of non-marine Molluscs from the Alburni Mountains, Campania, Southern Italy (Mollusca Gastropoda Bivalvia)

Agnese Petraccioli¹, Paolo Crovato², Ivano Niero³, Laura De Riso⁴, Camillo Pignataro⁵, Gaetano Odierna¹ & Nicola Maio^{1*}

¹Dipartimento di Biologia, Complesso Universitario di Monte S. Angelo, Università degli Studi di Napoli Federico II, Edificio 7, via Cinthia 21, 80126 Napoli, Italy

²Via S. Liborio 1, 80134 Napoli, Italy

³Via Cici 17/1, 30038 Spinea (Venezia), Italy

⁴Ente Parco Nazionale del Cilento, Vallo di Diano e Alburni, Via Montesani, 84078 Vallo Della Lucania (Salerno), Italy

⁵Fondazione I.RI.DI.A. - Museo Naturalistico, via Forese 16, 84020 Corleto Monforte (Salerno), Italy

*Corresponding author, e-mail: nicomaio@unina.it

ABSTRACT

An annotated checklist of the species of non-marine molluscs from the Alburni Mountains (Salerno Province, Campania, Southern Italy) is reported. The research was carried out from 2010 to 2013 inside a Site of Community Importance (SCI) and a Special Protection Area (SPA), of the Cilento, Vallo di Diano and Alburni National Park. The non-marine molluscs sampled on the field were compared with data available from the literature and malacological collections. Up to now, only 12 non-marine Mollusc species were known from the Alburni Mountains through bibliographical data. In all, the malacofauna of Alburni Mountains is composed by 83 non-marine Mollusc species (73 species of land snails, and 10 species of freshwater molluscs). The presence of nine species (six species of land snails and three species of freshwater snails) was confirmed by our field investigation, four species (3 species of land snails and 1 species of allochthonous freshwater snails) were recorded only by bibliographical data and were not yet found. Our analysis identifies 70 species of non-marine Molluscs (64 species of land snails, 6 species of freshwater molluscs) recorded on the basis of field data which were not previously recorded from the study area. At least 11 species are new records for the Campania Region. Extremely interesting is the record of *Vertigo angustior* Jeffreys, 1830 a species protected in European Union by the Annex II of the “Habitats Directive” and listed as “Vulnerable” at the European level. A Red List of Threatened Species is proposed and the species were classified with the code of I.U.C.N. (Version 2014.3). Five allochthonous species were surveyed for the first time in the study area: 3 land snails: *Lucilla scintilla* (Lowe, 1852), *Lucilla singleyana* (Pilsbry, 1829) and *Paralaoma servilis* (Shuttleworth, 1852), and 2 freshwater snails: *Potamopyrgus anti-podarum* (J.E. Gray, 1843) and *Ferrissia fragilis* (Tryon, 1863). Four species are known exclusively from the literature: *Vertigo (Vertigo) moulinsiana* (Dupuy, 1849), *Macrogastra (Pyrostoma) plicatula* (Draparnaud, 1801), *Cerņuella virgata* (Da Costa, 1778), and *Haitia acuta* (Draparnaud, 1805).

KEY WORDS

Non-marine Molluscs; Alburni Mountains, Campania; faunistics; conservation.

Received 18.07.2014; accepted 29.11.2014; printed 30.03.2015

Proceedings of the 2nd International Congress “Speciation and Taxonomy”, May 16th-18th 2014, Cefalù-Castelbuono (Italy)

INTRODUCTION

The knowledge of the malacofauna of Campania (about 150 species, personal data; 13,595 sq Km) is far below than that of other regions. This is even more evident when we consider the protect areas of this territory as those of the Alburni Mountains whose malacofauna is virtually unknown (Fig. 1).

In all Campania Region, only papers on checklist of Capri Island and Vesuvius National Park are known in the last 10 years (Petraccioli et al., 2005a, 2005b, 2006a, 2006b, 2007; Picariello et al., 2011); moreover there are only historical reports, nearly 100 years old, concerning the most common species and quoted with obsolete names. The purpose of this paper is, therefore, to help bridge this gap by increasing the malacofauna knowledge of this group in an important area of the Campania Region as the Alburni Mountains, with particular reference to the species listed in the Annex of the Habitats Directive, in the Index of the protected fauna of Italy and in the various red lists (Manganelli et al., 2000a; Cerfolli et al., 2002; I.U.C.N., 2014).

MATERIAL AND METHODS

The Alburni Mountains are a calcareous massif in the Salerno Province (Campania Region) belonging to the Lucan Sub Apennines chain, located in the Eastern area of Cilento, near the borders between Campania and Basilicata. In North-East the range degrades into the plain of Vallo di Diano between the valleys of the Calore Lucano, Tanagro and Sele rivers. The massif extends for about 250 km². The study area (SCI IT8050033 named: "Monti Alburni", and SPA IT8050055 named: "Alburni") is included in Cilento, Vallo di Diano and Alburni National Park and covers 14 administrative municipalities (Aqura, Auletta, Castelcivita, Controne, Corleto Monforte, Ottati, Petina, Polla, Postiglione, San Pietro al Tanagro, San Rufo, Sant'Angelo a Fasanella, Sant'Arzenio, and Sicignano degli Alburni) (see TEMI, 2010).

A detailed investigation on the historical and current literature and a comprehensive study of Neapolitan public and private molluscan collections were preliminarily performed. We also inspected the original sites reached by Costa (1874) 140 years ago.

From 2010 to 2013 the field surveys were conducted in 127 sampling points (stations or plots) between 100 and 1742 meters above sea level (sum-

mit of Monte Alburno/Panormo) in all suitable habitats present on the territory of the Alburni Mountains in accordance with the vegetation types reported in the land use map (1: 25.000) available by the "Ente Parco". Adult specimens and shells of non-marine molluscs were hand-collected through visual search, leaf litter and soil collecting and sorting. Samples were then air dried and sieved down to 0.5 mm mesh. Samples of sediment were screened with calibrated sieves. The cleaned up material was examined under lens and/or stereo microscope to sort the smallest fraction, namely. Fractions above 1 cm were searched by a Leica EZ4 stereo microscope (Leica Microsystems GmbH, Wetzlar, Germany), both incident and transmitted, and then photographed with a digital camera. The specimens for anatomical exams were drowned in water and fixed in 75% ethanol. The reproductive apparatus was extracted by means of scalpel, scissors and forceps. The illustrations of genitalia were sketched using a camera lucida mounted on the above stereomicroscope.

The sampled specimens were collected with permission of the "Ente Parco Nazionale del Cilento e Vallo di Diano" (Permit no. 16341/19.10.2010). Two sampled specimens for species were deposited in the Museo Naturalistico of Corleto Monforte (Salerno Province), a museum acknowledged as an "institution of regional interest" (Decreto dalla Giunta Regionale Campania n. 2010 del 29/12/2008). In addition, when other specimen/species were collected, we preserved them in the private collection of the authors. The species identification was based on qualified dichotomickeys (Giusti & Pezzoli, 1980; Girod et al., 1980; Bech, 1990; Giusti et al., 1995; Kerney & Cameron, 1999). The taxonomic order and nomenclatural arrangement of the list follow: Bodon et al. (1995), Manganelli et al. (1995, 1998, 2000b), Castagnolo (1995), Ponder & Lindberg (1996), Nordsieck (2002) and Bank (2011); the common names were based on Janus (1982) and on the web site: <http://media.eol.org>.

For each species a brief note on the abundance in the study area is reported according to the following classification: Very rare (sampled in 1–5 stations), Rare (sampled in 6–10 stations), Uncommon (sampled in 11–19 stations), Common (sampled in 20–35 stations), Widespread (sampled in over 35 stations). The bibliographical and museological data were then reported. If present in the I.U.C.N. Red List, each species is classified

with the code of Red List of Threatened Species (I.U.C.N., 2014) and Cuttelod et al. (2011).

data from 1986. Classified as “LC” by Cuttelod et al. (2011) and by I.U.C.N. (2014).

RESULTS

Species surveyed on the field

Phylum MOLLUSCA Cuvier, 1795
 Classis GASTROPODA Cuvier, 1795
 Subclassis ORTHOGASTROPODA Ponder et Lindberg, 1996

Ordo ARCHITAENIOGLOSSA Haller, 1890

Familia COCHLOSTOMATIDAE Kobelt, 1902

Cochlostoma montanum (Issel, 1866)
Cochlostoma montanum cassiniacum (Saint-Simon, 1878)

Common, 34 plots, locally abundant. Museal

Familia ACICULIDAE J.E. Gray, 1850

Platyla talentii Bodon et Cianfanelli, 2008

Rare, 10 plots, locally abundant (Figs. 2, 3). Endemic of Southern Apennine. Bodon & Cianfanelli, 2008. Classified as “NT” by Cuttelod et al. (2011) and “NT” by I.U.C.N. (2014).

Ordo NEOTAENIOGLOSSA Haller, 1892

Familia POMATIIDAE Newton, 1891 (1828)

Pomatias elegans (O.F. Müller, 1774)
 Round-mounted Snail

Common, 28 plots. Museal data from 1986.




Figure 1. The study area: Alburni Mountains, S-Italy.

Familia HYDROBIIDAE Stimpson, 1865
Mud snails

Pseudamnicola (P.) cfr. *moussonii* (Calcara, 1841)

Very rare, 1 plot. Classified as “LC” by Cuttelod et al. (2011) and by I.U.C.N. (2014). Endemic species in Europe (Cuttelod et al., 2011).

Belgrandia minuscula (Paulucci, 1881)

Very rare, 2 plots. Bodon et al. (2005). Classified as “DD” and Endemic species in Europe by Cuttelod et al. (2011) and “DD” by I.U.C.N. (2014).

Potamopyrgus antipodarum (J.E. Gray, 1843)
New Zealand mud snail, Jenkins' Spire Snail

Very rare, 2 plots (Fig. 4). Allochthonous species, introduced from New Zealand (Lori et al., 2005; Lori & Cianfanelli, 2007; Cianfanelli, 2009; Cuttelod et al., 2011). Classified as “LC” by I.U.C.N. (2014).

Bythinella opaca (M. von Gallenstein, 1848)
Bythinella schmidtii (Küster, 1852)

Very rare, 4 plots. Bodon et al. (1999) in an adjacent locality; Bodon et al., 2005. Classified as “LC” by Cuttelod et al. (2011) and by I.U.C.N. (2014). Endemic species in Europe (Cuttelod et al., 2011).

Familia ELLOBIIDAE Pfeiffer, 1854

Carychium tridentatum (Risso, 1826)
Long-toothed Herald Snail

Uncommon, 14 plots, locally abundant.

Familia LYMNAEIDAE Rafinesque, 1815

Galba truncatula (O.F. Müller, 1774)
Dwarf Pond Snail

Very rare, 2 plots. Museal data from 1986. Classified as “LC” by Cuttelod et al. (2011) and by I.U.C.N. (2014).

Radix labiata (Rossmässler, 1835)
Lymnaea (Radix) peregra (O.F. Müller, 1774)
Radix peregra (O. F. Müller, 1774)
Wandering Snail

Very rare, 3 plots. Costa (1874): sub *Limnaeus Gibilmannicus* (see O. G. Costa, 1839). Museal data from 1986. *Radix peregra* is classified by I.U.C.N. (2014) as synonym of *Radix balthica* (Linnaeus, 1758). *Radix labiata* is classified as “LC” by Cuttelod et al. (2011).

Ordo PULMONATA Cuvier in Blainville, 1814
Subordo BASOMMATOPHORA Keferstein, 1864
Familia ANCYLIDAE

Ancylus fluviatilis O.F. Müller, 1774
River Limpet

Rare, 7 plots. Classified as “LC” by Cuttelod et al. (2011) and by I.U.C.N. (2014).

Ferrissia fragilis (Tryon, 1863)
Ferrissia wautieri (Mirolli, 1960)
Fragile Ancyloid

Very rare, 1 plot (Fig. 5). Classified as “LC” by I.U.C.N. (2014). Cryptic invader of Italian freshwater ecosystems from North America (Cianfanelli et al., 2007; Lori & Cianfanelli, 2007).

Subordo STYLOMMATOPHORA A. Schmidt, 1855
Familia PYRAMIDULIDAE Kennard et B.B. Woodward, 1914

Pyramidula pusilla (Vallot, 1801)
Rock Snail

Common, 24 plots, locally abundant.

Pyramidula rupestris (Draparnaud, 1801)
Rock Snail

Very rare, 1 plot. Costa (1874).

Familia VERTIGINIDAE Fitzinger, 1833

Vertigo (Vertigo) pygmaea (Draparnaud, 1801)
Common Whorl Snail, Crested vertigo

Very rare, 1 plot. Classified as “LC” by Cuttelod et al. (2011)

Vertigo (Vertilla) angustior Jeffreys, 1830
Vertigo sinistrorso minore, Narrow-mouthed Whorl Snail

Very rare, 1 plot (Fig. 6). Species protected in European Union by the Annex II of the “Habitats Directive”, and in Italy by the D.P.R. n. 357/1997 than modified by D.P.R. n. 120/2003. In Europe, this species is listed as Vulnerable (VU) (criteria: A2ac+3c) at the European level and at the level of the 27 member States of the European Union (Cuttelod et al., 2011). The species is regionally protected in Tuscany, Umbria and Emilia-Romagna. The species has been regarded in Italy as “NT” by Manganelli et al. (2000a). Classified as “LR” by Cerfolli et al. (2002), and “NT” by I.U.C.N. (2014).

Columella edentula (Draparnaud, 1805)
Toothless Chrysalis Snail

Very rare, 2 plots. Classified as “LC” by Cuttelod et al. (2011) and by I.U.C.N. (2014). Endemic species in Europe (Cuttelod et al., 2011).

Truncatellina callicratis (Scacchi, 1833)

Rare, 10 plots. Endemic species in Europe classified as “LC” by Cuttelod et al. (2011).

Familia ORCULIDAE Pilsbry, 1918

Sphyradium doliolum (Bruguière, 1792)

Uncommon, 12 plots. Classified as “LC” by Cuttelod et al. (2011).

Pagodulina pagodula (des Moulins, 1830)

Very rare, 1 plot. Classified as “LC” by Cuttelod et al. (2011) and by I.U.C.N. (2014). Endemic species in Europe (Cuttelod et al., 2011).

Familia CHONDRINIDAE Steenberg, 1925

Rupestrella philippii (Cantraine, 1840)

Very rare, 4 plots. Classified as “LC” by Cuttelod et al. (2011).

Chondrina avenacea (Bruguière, 1792)

Widespread, 36 plots. Museal data from 1874. (Costa, 1874). Classified as “LC” by Cuttelod et al. (2011) and by I.U.C.N. (2014). Endemic species in Europe (Cuttelod et al., 2011).

Familia LAURIIDAE Steenberg, 1925

Lauria sempronii (Charpentier, 1837)

Rare, 8 plots.

Familia ARGNIDAE Hudec, 1965

Argna biplicata (Michaud, 1831)

Very rare, 4 plots. Classified as “LC” by Cuttelod et al. (2011) and by I.U.C.N. (2014). Endemic species in Europe (Cuttelod et al., 2011).

Familia VALLONIIDAE Morse, 1864

Acanthinula aculeata (O.F. Müller, 1774)
Pricly Snail

Common, 21 plots. Classified as “LC” by Cuttelod et al. (2011).

Gittenbergia sororcula (Benoit, 1859)

Rare, 9 plots. Locally very abundant. Classified as “LC” by Cuttelod et al. (2011).

Familia ENIDAE B.B. Woodward, 1903 (1880)

Chondrula tridens (O.F. Müller, 1774)

Very rare, 2 plots. Classified as “NT” by Cuttelod et al. (2011).

Jamina quadridens (O.F. Müller, 1774)

Uncommon, 17 plots. Locally very abundant. Museal data from 1986. Classified as “LC” by Cuttelod et al. (2011) and by I.U.C.N. (2014).

Merdigera obscura (O.F. Müller, 1774)

Ena obscura (O.F. Müller, 1774)

Lesser Bulin

Uncommon, 11 plots. Classified as “LC” by Cuttelod et al. (2011) and by I.U.C.N. (2014). Endemic species in Europe (Cuttelod et al., 2011).

Familia PUNCTIDAE Morse, 1864

Punctum pygmaeum (Draparnaud, 1801)

Dwarf Snail

Common, 33 plots.

Paralaoma servilis (Shuttleworth, 1852)

Paralaoma caputspinulae (Reeve, 1852)

Pinhead Spot

Very rare, 2 plots. Allochthonous species, introduced from New Zealand (Lori et al., 2005; Lori & Cianfanelli, 2007; Cianfanelli, 2009; Christensen, 2012).

Familia DISCIDAE Thiele, 1931 (1866)

Discus rotundatus (Müller, 1774)

Discus Snail, Rounded Snail

Rare, 7 plots.

Familia HELICODISCIDAE H.B. Baker, 1927

Lucilla scintilla (Lowe, 1852)

Oldfield Coil

Very rare, 1 plot (Fig. 7). Allochthonous species. The indigenous distribution for this species includes North America (Lori et al., 2005; Lori & Cianfanelli, 2007; Cianfanelli, 2009).

Lucilla singleyana (Pilsbry, 1889)

Smooth Coil

Very rare, 1 plot (Fig. 8). Allochthonous species. Originally probably from North America, introduced to Europe (Lori et al., 2005; Lori & Cianfanelli, 2007; Cianfanelli, 2009).

Familia VITRINIDAE Fitzinger, 1833

Vitrina* cfr. *pellucida (O.F. Müller, 1774)

Pellucid Glass Snail

Uncommon, 11 plots. Classified as “LC” by Cuttelod et al. (2011).

Familia PRISTILOMATIDAE T. Cockerell, 1891

Vitrea subrimata (Reinhardt, 1871)

Common, 26 plots.

Vitrea etrusca (Paulucci, 1878)

Very rare, 4 plots.

Vitrea contracta (Westerlund, 1871)

Milky Crystal Snail

Common, 26 plots.

Familia ZONITIDAE Mörch, 1864

Aegopis verticillus (Férussac, 1822)

Very rare, 1 plot (Fig. 9).

Familia OXYCHILIDAE P. Hesse, 1927 (1879)

Retinella olivetorum (Gmelin, 1791)

Retinella olivetorum olivetorum (Gmelin, 1791)

Common, 33 plots. Museal data from 1986.

Oxychilus* (*Oxychilus*) cfr. *draparnaudi (Beck, 1837)
Draparnaud's Glass Snail, Dark-bodied Glass snail

Uncommon, 16 plots. Museal data from 1986.

<i>Mediterranea hydatina</i> (Rossmässler, 1838)	Familia EUCONULIDAE Baker, 1928
Very rare, 1 plot.	
<i>Daudebardia rufa</i> (Draparnaud, 1805)	<i>Euconulus fulvus</i> (O.F. Müller, 1774)
Widespread, 42 plots. Locally very abundant (Figs. 10, 11, 12, 13).	Tawny Glass Snail, Brown Hive
	Rare, 8 plots.
Familia MILACIDAE Ellis, 1926	Familia FERUSSACIIDAE Bourguignat, 1883
<i>Tandonia sowerbyi</i> (A. Férussac, 1823)	<i>Cecilioides acicula</i> (O.F. Müller, 1774)
Keeled Slug, Sowerby's Slug	Blind Snail
Uncommon, 11 plots.	Very rare, 5 plots.
	<i>Cecilioides (Cecilioides) veneta</i> (Strobel, 1855)
Familia LIMACIDAE Lamarck, 1801	<i>Cecilioides janii</i> (De Betta et Martinati, 1855)
<i>Limax maximus</i> Linnaeus, 1758	Very rare, 5 plots.
Leopard Slug, Great Grey Slug, Giant Garde Slug	
Rare, 6 plots.	Familia SUBULINIDAE P. Fischer et Crosse, 1877
<i>Lehmannia marginata</i> (O. F. Müller, 1774)	<i>Rumina decollata</i> (Linnaeus, 1758)
Tree slug	Decollate Snail
Very rare, 1 plot.	Very rare, 4 plots.
<i>Limacus flavus</i> (Linnaeus, 1758)	Familia OLEACINIDAE H. Adams et A. Adams, 1855
Yellow Slug, Tawny Garden Slug	<i>Poiretia dilatata</i> (Philippi, 1836)
Very rare, 2 plots.	Common, 31 plots. Museal data from 1986.
Familia AGRIOLIMACIDAE H. Wagner, 1935	Familia TESTACELLIDAE J.E. Gray, 1840
<i>Deroceras reticulatum</i> (O.F. Müller, 1774)	<i>Testacella scutulum</i> G.B. Sowerby I, 1821
Netted Slug, Gray Fieldslug	Shield Shelled Slug
Rare, 6 plots.	Very rare, 4 plots.
<i>Deroceras invadens</i> Reise, Hutchinson, Schun- ach et Schlitt, 2011	Familia CLAUSILIIDAE J.E. Gray, 1855
Chestnut Slug, Brown Field Slug, Longneck Fieldslug, Widespread Pest Slug	Door Snails
Very rare, 4 plots.	<i>Medora</i> sp.
	Very rare, 3 plots.

Cochlodina (Cochlodina) laminata (Montagu, 1803)
Plaited Door Snail

Rare, 6 plots. (Costa, 1874).

Charpentieria (Stigmatica) paestana (Philippi, 1836)

Siciliaria paestana (Philippi, 1836)

Widespread, 42 plots. Locally abundant.

Charpentieria (Stigmatica) cfr. ernae (Fauer, 1978)

Very rare, 2 plots. Museal data from 1970. Endemic of Southern Appennine. (Fauer, 1978; Welter-Schultes, 2012; Nordsieck, 2013).

Macrogastra (Pyrostoma) attenuata (Rossmässler, 1835)

Macrogastra (Pyrostoma) attenuata iriana (Pol-lonera, 1885)

Lined door snail

Very rare, 2 plots.

Clausilia cruciata (S. Studer, 1820)

Clausilia cruciata bonellii E. Von Martens, 1873

Very rare, 3 plots. Museal data before 1950.

Familia HYGROMIIDAE Tryon, 1866

Xerotricha conspurcata (Draparnaud, 1801)

Very rare, 2 plots. Museal data from 1986. Classified as “LC” by Cuttelod et al. (2011).

Hygromia cinctella (Draparnaud, 1801)

Girdled Snail

Rare, 8 plots. Classified as “LC” by Cuttelod et al. (2011) and by I.U.C.N. (2014). Endemic species in Europe (Cuttelod et al., 2011).

Cerneuella (Cerneuella) cisalpina (Rossmässler, 1837)

Common, 21 plots. Locally abundant. Museal data from 1986. Classified as “LC” by Cuttelod et al. (2011) and by I.U.C.N. (2014). Endemic species in Europe (Cuttelod et al., 2011).

Cerneuella (Xerocincta) neglecta (Draparnaud, 1805)
Luddesdown Snail, Neglected dune snail

Very rare, 3 plots. Museal data from 1986. Classified as “LC” by Cuttelod et al. (2011) and by I.U.C.N. (2014). Endemic species in Europe (Cuttelod et al., 2011).

Cerneuellopsis ghisottii Manganelli et Giusti, 1988

Very rare, 3 plots. Hallgas, com. pers., 2013. This species is endemic to Italy. Classified as “VU” by Cuttelod et al. (2011) and by I.U.C.N. (2014) (IUCN Criteria (Europe) (version 3.1): B1ab(iii)+2ab(iii). Endemic species in Europe (Cuttelod et al., 2011).

Trochoidea (Trochoidea) pyramidata (Draparnaud, 1805)

Very rare, 2 plots. Classified as “LC” by Cuttelod et al. (2011).

Trochoidea (Trochoidea) trochoides (Poiret, 1789)

Very rare, 2 plots. Museal data from 1985. Classified as “LC” by Cuttelod et al. (2011).

Monacha (Monacha) cfr. cartusiana (O.F. Müller, 1774)

Chartreuse Snail, Carthusian Snail


Common, 35 plots. Museal data from 1986. Classified as “LC” by Cuttelod et al. (2011) and by I.U.C.N. (2014). Endemic species in Europe (Cuttelod et al., 2011). Regarded as “Edible species” in Tuscany and Umbria Region.

Monacha (Eutheba) cfr. parumcincta (Menke, 1828)

Common, 26 plots. Classified as “LC” by Cuttelod et al. (2011) and by I.U.C.N. (2014). Endemic species in Europe (Cuttelod et al., 2011). Familia HELICODONTIDAE Kobelt, 1904

Helicodonta obvoluta (O.F. Müller, 1774)

Helicodonta obvoluta obvoluta (O.F. Müller, 1774)
Cheese Snail



Figures 2-9. Non-marine Molluscs from the Alburni Mountains, Campania, Southern Italy. Figure 2. *Platyla talentii* alive (Photo by N. Maio). Figure 3. Shell of *Platyla talentii*: view. 3A. Particular of dorsal view. 3B. Particular of the mouth (Photos by N. Maio). Figure 4. *Potamopyrgus antipodarum* alive (Photo by N. Maio). Figure 5. Shell of *Ferrissia fragilis*: dorsal, lateral and ventral view (Photo by I. Niero). Figure 6. Shell of *Vertigo (Vertilla) angustior*: apertural view (Photo by I. Niero). Figure 7. Shell of *Lucilla scintilla*: umbilical, dorsal and apertural view (Photo by N. Maio). Figure 8. Shell of *Lucilla singleyana*: dorsal, apertural and umbilical view (Photo by N. Maio). Figure 9. Shell of *Aegopis verticillus* dorsal, umbilical and apertural view (Photo by N. Maio).

Rare, 7 plots. Classified as “LC” by Cuttelod et al. (2011) and by I.U.C.N. (2014). Endemic species in Europe (Cuttelod et al., 2011).

Familia HELICIDAE Rafinesque, 1815

Chilostoma (Campylea) cfr. planospira (Lamarck, 1822)

Uncommon, 18 plots. Museal data from 1986. Classified as “DD” by Cuttelod et al. (2011) and by I.U.C.N. (2014). Endemic species in Europe (Cuttelod et al., 2011).

Marmorana (Ambigua) fuscolabiata fuscolabiata (Rossmässler, 1842)

Widespread, 27 plots. Costa (1874), Degner (1927). Museal data from 1874. Classified as “DD” by Cuttelod et al. (2011) and by I.U.C.N. (2014). Endemic species in Europe (Cuttelod et al., 2011).

Marmorana (Ambigua) cfr. fuscolabiata wullei Kobelt, 1903

Common, 25 plots. Kobelt (1903a; 1903b), Degner (1927), Bacci (1951), Alzona (1971). Museal data from 1903.

Eobania vermiculata (O.F. Müller, 1774)
Chocolate-band snail

Very rare, 3 plots. Museal data from 2006. Regarded as “Edible species” in Tuscany and Umbria Region.

Cantareus apertus (von Born, 1778)
Green Garden Snail

Very rare, 2 plots. Museal data from 1986. Regarded as “Edible species” in Tuscany and Umbria Region.

Cornu aspersum (O.F. Müller, 1774)
Cantareus aspersus (O.F. Müller, 1774)
Garden Snail, Common Snail, Brown Garden Snail

Very rare, 4 plots. Museal data from 1986. Regarded as “Edible species” in Tuscany and Umbria Region.

Helix (Helix) cfr. delpretiana Paulucci, 1878

Rare, 9 plots. Museal data from 1986. This species is endemic to the Central Appenines in Italy. Classified as “DD” by Cuttelod et al. (2011) and by I.U.C.N. (2014). Endemic species in Europe (Cuttelod et al., 2011).

Helix (Helix) cfr. ligata O.F. Müller, 1774
Ligate Snail

Uncommon, 11 plots. Museal data from 1986. Classified as “LC” by Cuttelod et al. (2011) and by I.U.C.N. (2014). Endemic species in Europe (Cuttelod et al., 2011).

Classis BIVALVIA Linnaeus, 1758

Ordo VENEROIDA H. et A. Adams, 1857
Familia SPHAERIIDAE Deshayes, 1855 (1820)

Pisidium casertanum (Poli, 1791)
Caserta Pea Mussel

Very rare, 5 plots. Classified as “LC” by Cuttelod et al. (2011) and by I.U.C.N. (2014).

Species exclusively known from the literature

Familia VERTIGINIDAE Fitzinger, 1833

Vertigo (Vertigo) moulinsiana (Dupuy, 1849)
Vertigo of Demoulins, Demoulins’ Whorl Snail

Manganelli et al. (2001), Bodon et al. (2005). Find only in debris of Fiume Calore, near Grotta di Castelcivita, Salerno Province, by S. Cianfanelli and E. Talenti on 1994. Species protect in European Union by the Annex II of the “Habitats Directive” and in Italy by the D.P.R. n. 357/1997 than modified by D.P.R. n. 120/2003. The species is regionally protect in Tuscany, Umbria and Emilia-Romagna. The species has been initially classified as “LRcd”(= Lower risk, conservation dependant) by Bouchet et al. (1999), then the species has been regarded in Italy as “VU”(criteria: B2a, B2b) by Manganelli et al. (2000b, 2001). Classified as “LR” by Cerfolli et al. (2002), “VU” (criteria: A2ac) by Cuttelod et al. (2011) and by I.U.C.N. (2014).




Figure 10. Shell of *Daudebardia rufa*: dorsal, apertural and umbilical view (Photo by I. Niero). Figure 11. *D. rufa* alive (Photo by N. Maio). Figure 12. Genitalia of *D. rufa* (Sant'Angelo a Fasanella (SA), 1160 m, 17.0V.2013, N. Maio, P. Crovato & I. Niero legit). Figure 13. Internal structure of the penis. Acronyms in figures: AG= albumen gland; DBC= duct of the bursa copulatrix; E= epiphallus; EP= epiphallic pore; FO= free oviduct; GA= genital atrium; HD= hermaphroditic duct; HG= hermaphroditic gland; P= penis; PPL= penis pleats; PR= penial retractor; Pro= prostate; PVG= perivaginal gland; PS = penis sheath; RS = reservoir of spermatheca; T= talon; UOS= uterine ovispermiduct; Va= vagina, VD= vas deferens. (Drawn by I. Niero).

Familia CLAUSILIIDAE J.E. Gray, 1855

Macrogastra cfr. *plicatula* (Draparnaud, 1801)

Costa (1874: sub *Clausilia plicatula* Drap.).
Museum data before 1950.

From the South of Italy are cited 3 subspecies, in addition to *M. plicatula plicatula*: *M. p. amiaten-sis* Nordsieck, 2006; *M. p. apennina* Gentiluomo 1868, and *M. p. aprutina* Nordsieck, 2006 (Nordsieck, 2006).

Familia HYGROMIIDAE Tryon, 1866

Cernuella cfr. *virgata* (Da Costa, 1778)

Banded Snail

Kobelt (1907). Museum data from 1986. Regarded as “Edible species” in Tuscany and Umbria Region.

Familia OXYCHILIDAE P. Hesse, 1927 (1879)

Oxychilus sp.

Capasso (1958), Capolongo & Cantilena (1974).

Ordo HYGROPHILA A. Férussac 1822

Familia PHYSIDAE Fitzinger, 1833

Haitia acuta (Draparnaud, 1805)

Acute Bladder Snail, European physa, Tadpole Snail, Bladder Snail, Pewter physa

Sacchi (1964): sub *Physa acuta*. Allochthonous species. The oldest alien species of Italy probably native to northeastern North America (Lori et al., 2005; Lori & Cianfanelli, 2007; Cianfanelli, 2009).

DISCUSSION

Up to now, only 13 non-marine Mollusc species were known from the Alburni Mountains through bibliographical data (nine species of land snails and four species of freshwater snails), to which we add the new species listed in this paper for the study

area. The presence of nine species (six species of land snails and three species of freshwater molluscs) are confirmed by our field investigation. Only four species (the land snails *Macrogastra plicatula*, *Vertigo moulinsiana*, *Cernuella virgata* and the allochthonous freshwater snails *Haitia acuta*) were documented exclusively by bibliographical data and have not been confirmed by the field surveys yet (Fiorentino et al., 2008; Reise et al., 2011). Our analysis identifies 79 species of non-marine molluscs (69 species of land snails, ten species of freshwater molluscs) recorded on the basis of field data. In total the occurrence of 83 species of non-marine molluscs (73 species of land snails, 10 species of freshwater molluscs) was herein attested in the survey area representing approximately the 56% of the estimated fauna of Campania Region (about 150 species, personal data).

At least 11 species are new records for the Campania Region (*Aegopsis verticillus*, *Cernuellopsis ghisottii*, *Lucilla scintilla*, *Argna biplicata*, *Cernuella neglecta*, *Daudebardia rufa*, *Helix delpretiana*, *Pagodulina pagodula*, *Vitrina pellucida*, *Vitrea etrusca* and *Macrogastra attenuata*). *M. attenuata* (sub *M. lineolata*) and *D. rufa* were generically recorded from Matese Mountains (probably Molise) by Giusti et al. (1985). 70 species of non-marine molluscs (64 species of land snails, six species of freshwater molluscs), recorded on the basis of field data, have not been previously recorded from the study area.

Extremely interesting is the finding of samples of *Medora* sp.: it seems to be the second record of this genus for the region. The systematics of the genus *Medora* Adams, 1855 is in fact complex and, in many respects, still controversial. Regarding Italy, Nordsieck (1970) considered *M. italiana* (Küster, 1847) of the Central-Southern Apennines distinct from *M. albescens* (Menke, 1830) of the Balkan peninsula. In addition, he assigned to *M. italiana* various subspecies: only one from Campania: *M. i. italiana* (Küster, 1847) (locus typicus: Piedimonte d'Alife (= Piedimonte Matese, Caserta, Campania). Giusti et al. (1986) suggested that it was not possible to distinguish *M. italiana* from *M. albescens* with the subspecies *M. a. italiana* in the central part of Italy. The populations reported for Italy as *M. dalmatina* (Manganelli et al., 1995) were described by Nordsieck (2012) as a distinct sub-

species. Preliminary data by Colomba et al. (2012) suggest that the genus *Medora* shows a much more complex and articulate differentiation than hitherto hypothesized by morphological surveys so far. An attempt to clarify its organization and internal structure, at various taxonomic levels, a more detailed analysis including a higher number of molecular markers and additional *Medora* populations from Italy are required.

Other interesting records are: *Vertigo angustior*, a species protected in the European Union by the Annex II of the Council Directive 92/43/EEC of May 21st 1992 on the conservation of natural habitats and of wild fauna and flora known as “Habitats Directive”, that includes “animal and plant species of community interest whose conservation requires the designation of special areas of conservation (SPA)” and listed as “Vulnerable” at the European level and *Platyla talentii*, an endemic species of Southern Apennine, recently described by Bodon & Cianfanelli (2008), classified as “Near Threatened” by Cuttelod et al. (2011) and by I.U.C.N. (2014).

Five allochthonous species were surveyed for the first time in the study area: three land snails (*Lucilla scintilla*, *L. singleyana* and *Paralaoma servilis*) and two freshwater snails (*Potamopyrgus antipodarum* and *Ferrissia fragilis*). *L. scintilla* and *L. singleyana* are native in North America; they were probably introduced into Europe in the second half of the 20th century (Horsák et al., 2009). *L. singleyana*, *P. servilis* and *F. fragilis* are the second records for the Campania Region, the first is Bodon et al. (2004) sub *Helicodiscus singleyanus* (Pilsbry, 1890), Bodon et al. (2004) sub *P. caputspinulae* (Reeve, 1852) and D'Antonio & Bravi (1990) sub *F. wautieri* (Mirolli, 1960).

In the past, three endemic taxa had been described as new for the Alburni area: *Helix (Iberus) wullei* first described by Kobelt (1903a: 14-15, tavv. 1766-1768) from “Monte Alburno circa vicum Postiglione prov. Salernitanae” and later as *Iberus wullei* (Kobelt, 1903b: pag. 4-5, fig. without number); *Xerophila (Xerolauta) peninsularis* forma “alburni” described by Kobelt (1907: pp. 59-60, Tafel 357 Fig. 2221) from “Monte Postiglione, des alten Alburnus” and *Siciliaria ernae* described by Fauer (1978: 265, abb. 1) from “Passo Sentinella im SO der Monti Aburni” [municipality of Corleto Monforte], and “6 km West of San Rufo”. Today *X.*

peninsularis is considered a junior synonym of *Cernuella virgata* (Da Costa, 1778) but the taxonomic status of *Iberus* (= *Marmorana*) *wullei* and of *S. ernae* need to be confirmed. The Demoulin's Whorl Snail, listed in the Standard Data Form of the SCI of “Monti Alburni” is not confirmed by our field surveys.

A Red List of Threatened Species is also proposed and the species were classified with the code of I.U.C.N. (Version 2014.3).

ACKNOWLEDGEMENTS

This research was funded with the support of the Cilento, Vallo di Diano and Alburni National Park. Authors wish to thank: the National Park Authority of the Cilento, Vallo di Diano and Alburni for collaboration as well as advising on technical and administrative aspects. In particular: Amilcare Troiano (President), Corrado Matera (Vice-President) and Angelo De Vita (Director). A special thank goes to Pasquale and Pietro Cappelli (Ottati, Salerno) for the trekking guide. We want to thank: Mario Cuomo, Sergio Duraccio, Gianni D'Anna, Franco Izzillo, and Massimo Cretella (Naples) and Alessandro Hallgass (Rome) for the unpublished data given; Antonio Pietro Ariani for the malacological collection consultation of the Zoological Museum of Naples; Monica Leonardi (Conservator) for the malacological collection consultation of the Museo Civico di Storia Naturale di Milano, Maria Tavano (Conservator) and Giacomo Doria (Director) for the malacological collection consultation of the Museo Civico di Storia Naturale di Genova “Giacomo Doria”. Furthermore we thank: Domenico Davolos (Dipartimento di Biologia, Università di Napoli Federico II), Antonietta Mordente (Banca Monte Pruno, Credito Cooperativo di Roscigno e Laurino) and Nicola Auricchio (President of Fondazione I.R.I.D.I.A. - Museo Naturalistico of Corleto Monforte) and Antonio Sicilia (Mayor of Municipality of Corleto Monforte) for administrative support and cooperation.

REFERENCES

Alzona C., 1971. Malacofauna italiana. Catalogo e bibliografia dei molluschi viventi, terrestri e d'acqua dolce.

- Parte I. Atti della Società Italiana di Scienze Naturali e del Museo Civico di Storia Naturale di Milano, Milano, 111: 1–433.
- Bacci G., 1951. Le razze di *Ambigua fuscolabiata* (Rossm.) (Polmonata - Helicidae): un problema di sistematica e di genetica. *Annuario dell'Istituto e Museo di Zoologia dell'Università di Napoli* (N. S.), 3: 10–24.
- Bank R. A., 2011. Fauna Europaea: Mollusca Gastropoda. Fauna Europaea version 2.4. Checklist of the land and freshwater Gastropoda of Italy. Fauna Europaea Project: 1-49. http://www.nmbe.unibe.ch/sites/default/files/uploads/pubinv/fauna_europaea_-_gastropoda_of_italy.pdf. Last update: November 24th, 2013.
- Bech M., 1990. Fauna malacològica de Catalunya. Molluscs terrestres i d'aigua dolça. Treballs de la Institució Catalana d'Història Natural. N. 12. Barcelona.
- Bodon M. & Cianfanelli S., 2008. Una nuova specie di *Platyla* per il sud Italia (Gastropoda: Prosobranchia: Aciculidae). *Bollettino Malacologico*, 44: 27–37.
- Bodon M., Favilli L., Giannuzzi Savelli R., Giovine F., Giusti F., Manganelli G., Melone G., Oliverio M., Sabelli B. & Spada G., 1995. Gastropoda; Prosobranchia, Heterobranchia Heterostropha. In: Minelli A., Ruffo S. & La Posta S. (Eds.), Checklist delle specie della fauna italiana. Calderini, Bologna, 14, 60 pp.
- Bodon M., Cianfanelli S., Talenti E., Manganelli G. & Giusti F., 1999. *Litthabitella chilodia* (Westerlund, 1886) in Italy (Gastropoda Prosobranchia: Hydrobiidae). *Hydrobiologia*, 411: 175–189.
- Bodon M., Lori E. & Cianfanelli S., 2004. Un'altra specie aliena per la malacofauna italiana: *Hawaia minuscula* (Binney, 1840) (Pulmonata: Zonitidae). *Bollettino Malacologico*, 40: 11–14.
- Bodon M., Cianfanelli S., Manganelli G., Pezzoli E. & Giusti F., 2005. Gastropoda Prosobranchia e Heterobranchia Heterostropha d'acqua dolce. In: Ruffo S. & Stoch F. (Eds.), Checklist e distribuzione della fauna italiana. Memorie del Museo Civico di Storia naturale di Verona, 2 Serie Sez. Scienze della Vita, 16: 79-81 + Cdrom. <http://ckmap.faunaitalia.it>.
- Bouchet P., Falkner G., Seddon M. B., 1999. Lists of protected land and freshwater molluscs in the Bern Convention and European Habitats Directive: are they relevant to conservation? *Biological Conservation*, 90: 21–31.
- Castagnolo L., 1995. Bivalvia (specie d'acqua dolce: generi 064-066, 128, 132-134). In: Minelli A., Ruffo S. & La Posta S. [a cura di], Checklist delle specie della fauna d'Italia. Calderini, Bologna. 17 (Bivalvia, Scaphopoda), 21 pp.
- Capasso P., 1958. Speleologia termale. Nota preliminare sulla località di Contursi (Salerno) e le sue mofete. *Studia Spelaologica*, 3 (giugno): 95–102.
- Capolongo D. & Cantilena S., 1974. Specie cavernicole di Campania. *Annuario dell'Istituto e Museo di Zoologia dell'Università di Napoli*, 20: 1–198.
- Cerfolli F., Petrassi F. & Petretti F., 2002. Il Libro Rosso degli Animali Invertebrati d'Italia. WWF Italia - ONLUS – Roma.
- Christensen C., 2012. First Records of *Paralaoma servilis* (Shuttleworth, 1852) (Gastropoda: Pulmonata: Punctidae) in the Hawaiian Islands. *Records of the Hawaii Biological Survey for 2011*. Bishop Museum Occasional Papers, 112: 3–7.
- Cianfanelli S., 2009. I Molluschi della Provincia di Pistoia: le specie da tutelare e quelle da combattere. *Quaderni del Padule di Fucecchio* n. 6. Centro di Ricerca, Documentazione e Promozione del Padule di Fucecchio, 112 pp.
- Cianfanelli S., Lori E. & Bodon M., 2007. Non-indigenous Alien freshwater molluscs in Italy and their distribution. In: Gherardi F. (Ed.). *Biological invaders in inland waters: profiles, distribution, and threats*. Springer, Dordrecht, The Netherlands. Chapter 5, pp 103-121, http://dx.doi.org/10.1007/978-1-4020-6029-8_5
- Colomba M.S., Liberto F., Reitano A., Renda W., Pocaterra G., Gregorini A. & Sparacio I., 2012. Molecular studies on the genus *Medora* H. et A. Adams, 1855 from Italy (Gastropoda Pulmonata Clausiliidae). *Biodiversity Journal*, 3: 571–582.
- Costa O. G., 1839. Cenni sulla Fauna Siciliana. In: Costa O. G. (a cura di), *Corrispondenza zoologica destinata a diffondere nel Regno delle Due Sicilie tutto ciò che si va discoprendo entro e fuori Europa (e vice-versa) riguardante la zoologia in generale*. Tip. Azzolino & Compagno, Napoli. Anno I (11-12): 159–184.
- Costa A., 1874. Una peregrinazione zoologica su' monti dell'Alburno. *Rendiconti della Reale Accademia delle scienze fisiche e matematiche di Napoli* (serie 1), 13: 129–135.
- Cuttelod A., Seddon M. & Neubert E., 2011. *European Red List of Non-marine Molluscs*. Luxembourg: Publications Office of the European Union, 98 pp.
- D'Antonio C. & Bravi S., 1990. Ricerche naturalistiche nella tenuta Astroni (Napoli). I. La malacofauna acquatica. *Lavori S.I.M., Atti Congresso di Sorrento* 23–31 maggio 1987, 23: 483–493.
- Degner E., 1927. Zur Molluskenfauna Unteritaliens. *Mitteilungen aus dem Zoologischen Staatsinstitut und Zoologischen Museum in Hamburg*, 43: 19–114.
- Fauer W., 1978. *Siciliaria ernae* n. sp., eine neue rezente Clausilie aus Italien. *Archiv für Molluskenkunde*, 108: 263–265.
- Fiorentino V., Salomone N., Manganelli G. & Giusti F., 2008. Phylogeography and morphological variability in land snails: the Sicilian *Marmorana* (Pulmonata, Helicidae). *Biological Journal of the Linnean Society*, 94: 809–823.

- Girod A., Bianchi I. & Mariani M., 1980. Gasteropodi, 1 (Gastropoda: Pulmonata, Prosobranchia: Neritidae, Bithyniidae, Valvatidae). Guida per il riconoscimento delle specie animali delle acque interne italiane. C.N.R., Roma, 86 pp.
- Giusti F. & Pezzoli E., 1980. Gasteropodi, 2 (Gastropoda: Prosobranchia: Hydrobioidea, Pyrguloidea). Guida per il riconoscimento delle specie animali delle acque interne italiane. C.N.R., Roma, 67 pp.
- Giusti F., Grappelli C., Manganelli G., Fondi R. & Bulli L., 1986. An attempt of natural classification of the genus *Medora* in Italy and Yugoslavia, on the basis of conchological, anatomical and allozymic characters (Pulmonata: Clausiliidae). *Lavori della Società Italiana di Malacologia*, 22: 259–341.
- Giusti F., Castagnolo L. & Manganelli G. 1985. La fauna malacologica delle faggete Italiane: brevi cenni di ecologia, elenco delle specie e chiavi per il riconoscimento dei generi e delle entità più comuni. *Bollettino Malacologico*, 21: 69–144.
- Giusti F., Manganelli G. & Schembri P.J., 1995. The non-marine molluscs of the Maltese Island. Museo Regionale di Scienze Naturali, Torino, 607 pp.
- Horsák M., Šteffek J., Čejka T., Ložek V. & Juříčková L., 2009. Occurrence of *Lucilla scintilla* (R.T. Lowe, 1852) and *Lucilla singleyana* (Pilsbry, 1890) in the Czech and Slovak Republics - with remarks how to distinguish these two non-native minute snails. *Malacologica Bohemoslovaca*, 8: 24–27.
- I.U.C.N., 2014. IUCN Red List of Threatened Species. Version 2014.3. Disponibile su www.iucnredlist.org. Downloaded on 22 June 2014.
- Janus H., 1982. The illustrated Guide to Molluscs. Harold Starke Limited, London, 180 pp.
- Kerney M.P. & Cameron R.A. D., 1999. Guide des escargots et limaces d'Europe identification et biologie de plus de 300 espèces. Delachaux et Niestlé, Paris, 370 pp.
- Kobelt W., 1903a. N. 1766-1768. *Helix (Iberus) wullei* n. Iconographie der Land- und Süßwasser-Mollusken mit vorzüglicher Berücksichtigung der Europäischen noch nicht abgebildeten Arten von E. A. Rossmässler. Neue Folge, vol. 10. C.W. Kreidel's Verlag, Wiesbaden, 107 pp.
- Kobelt W., 1903b. Diagnoses Heliceorum novorum in Italia collectorum. *Annuario del Museo Zoologico della Reale Università di Napoli (N. S.)*, 1: 1–5.
- Kobelt W., 1907. Iconographie der Land- & Süßwasser-Mollusken mit vorzüglicher Berücksichtigung der Europäischen noch nicht abgebildeten Arten von E.A. Rossmässler. C.W. Kreidel's Verlag, Wiesbaden, n.f. [n.s.], tomo [band] 13, pp. 1–65, tavv. 331–360.
- Lori E. & Cianfanelli S., 2007. Studio sulla presenza e distribuzione di Molluschi terrestri e d'acqua dolce alieni nel territorio della Provincia di Pistoia. Relazione finale 2007. Museo di Storia Naturale dell'Università degli Studi di Firenze, 97 pp.
- Lori E., Bodon M. & Cianfanelli S., 2005. Molluschi continentali alieni in Italia: presenza e distribuzione. *Notiziario S.I.M.*, 23: 71.
- Manganelli G., Bodon M., Favilli L. & Giusti F., 1995. Gastropoda Pulmonata. In: Minelli A., Ruffo S., La Posta S. [a cura di], Checklist delle specie della fauna d'Italia, Bologna, 16, 60 pp.
- Manganelli G., Bodon M., Favilli L., Castagnolo L. & Giusti F., 1998. Checklist delle specie della fauna d'Italia, molluschi terrestri e d'acqua dolce. Errata ed addenda, 1. *Bollettino Malacologico*, 33: 151–156.
- Manganelli G., Bodon M. & Giusti F., 2000a. Checklist delle specie della fauna d'Italia, molluschi terrestri e d'acqua dolce. Errata e addenda, 2. *Bollettino Malacologico*, 36: 125–130.
- Manganelli G., Bodon M., Cianfanelli S., Favilli L., Talenti E. & Giusti F., 2000b. Conoscenza e conservazione dei molluschi non marini italiani: lo stato delle ricerche. *Bollettino Malacologico*, 36: 5–42.
- Manganelli G., Cianfanelli S., Brezzi M. & Favilli L., 2001. The distribution and Taxonomy of *Vertigo moulinsiana* (Dupuy, 1849) in Italy (Gastropoda: Pulmonata: Vertiginidae). *Journal of Conchology*, 37: 267–280.
- Nordsieck H. 1970. Zur Anatomie und Systematik der Clausilien, VIII. Dinarische Clausiliidae, II: Das Genus *Medora*. *Archiv für Molluskenkunde*, 100: 23–75.
- Nordsieck H., 2002. Contributions to the knowledge of the Delimini (Gastropoda): Stylommatophora: Clausiliidae. *Mitteilungen der Deutschen malakozoologischen Gesellschaft*, 67: 27–39.
- Nordsieck H., 2006. Systematics of genera *Macrogastra* Hartmann 1841 and *Julica* Nordsieck 1963, with the description of new taxa (Gastropoda: Stylommatophora: Clausiliidae). *Archiv für Molluskenkunde*, 135: 49–71.
- Nordsieck H., 2012. Ergänzung der Revision der Gattung *Medora* H. & A. ADAMS: Die *Medora*-Arten Italiens (Gastropoda, Stylommatophora, Clausiliidae, Alopinae), mit Beschreibung einer neuen Unterart von *Medora dalmatina* Rossmässler. *Conchyliologia*, 42: 75–81.
- Nordsieck H., 2013. Delimini (Gastropoda, Pulmonata, Clausiliidae) from Apennine Italy, with the description of three new subspecies from Calabria. *Conchyliologia*, 44: 3–14.
- Petraccioli A., Barattolo F., Crovato P., Cretella M., Maio N. & Aprea G., 2005a. Guida pratica al riconoscimento dei macro-gasteropodi terrestri attuali e fossili dell'Isola di Capri. *Bollettino della Sezione Campania ANISN (N. S.)*, 16 (29): 19–48.

- Petraccioli A., Crovato P., Cretella M., Maio N., Aprea G. & Barattolo F., 2005b. I Gasteropodi continentali recenti dell'Isola di Capri. In: Crovato P. & Maio N. (Eds.). IV International Congress of the European Malacological Societies. October 10–14 2005, Naples (Italy). Abstracts of Oral Communications and Posters. *Notiziario S.I.M.*, 23: 28.
- Petraccioli A., Crovato P., Cretella M., Maio N., Aprea G. & Barattolo F., 2006a. I Gasteropodi continentali dell'Isola di Capri: risultati preliminari. In: Guglielmi R. & Nappi A. (Eds.), *Atti del Convegno: "La Natura in Campania: aspetti biotici e abiotici"*. Napoli, 18 novembre 2004. Gruppo Attivo Campano A.R.C.A. - Onlus, pp. 77–86.
- Petraccioli A., Maio N., Crovato P. & Picariello O., 2006b. I Gasteropodi terrestri del Parco Nazionale del Vesuvio. Risultati preliminari. 67° Congresso Nazionale Unione Zoologica Italiana. Napoli, 12–15 settembre 2006. Riassunti dei contributi scientifici: 84–85.
- Petraccioli A., Crovato P., Cretella M., Maio N., Aprea G. & Barattolo F., 2007. The fossil land Gastropods from Capri Island. IV International Congress of the European Malacological Societies. October 10–14 2005, Naples (Italy). *Bollettino Malacologico*, 43: 51–56.
- Picariello O., Maio N., Petraccioli A., Crovato P. & Guarino F.M., 2011. Emergenze Faunistiche del Parco Nazionale del Vesuvio. Atti XV Convegno Nazionale A.N.I.S.N.: "La Natura, l'Uomo, il Tempo". Napoli, 7–12 settembre 2010. *Le Scienze Naturali nella Scuola. Periodico A.N.I.S.N.*, 20 (44–III): 81–93.
- Ponder W.F. & Lindberg D.R., 1996. Gastropod phylogeny: Challenges for the 90s. In: Taylor J. (Ed.), *Origin and evolutionary radiation of the Mollusca*. Oxford University press., Oxford, pp. 135–154.
- Reise H., Hutchinson J.M.C., Schunack S. & Schlitt B., 2011. *Deroceras panormitanum* and congeners from Malta and Sicily, with a redescription of the widespread pest slug as *Deroceras invadens* n. sp. *Folia Malacologica*, 19: 201–233.
- Sacchi C.F., 1964. Origini ed evoluzione della malacofauna appenninica meridionale. *Annuario dell'Istituto e Museo di Zoologia dell'Università di Napoli*, 15/1963 (7): 1–85.
- TEMI s.r.l. (a cura di) (2010). Piano di gestione del Sito di Importanza Comunitaria "Monti Alburni" (IT8050033) e della Zona di Protezione Speciale "Alburni" (IT805055). Progetto Life Natura "LIFE06NAT/IT/000053" "Gestione della Rete di SIC/ZPS nel PN del Cilento e Vallo di Diano" (Cilento in Rete), 294 pp.
- Welter-Schultes F.W., 2012. European non-marine molluscs, a guide for species identification. Planet Poster Editions, Göttingen, 679 pp.