

RESEARCH ARTICLE

Open Access

Surgical mistake causing an high recto-vaginal fistula. A case report with combined surgical and endoscopic approach: therapeutic considerations

Michele Danzi^{1*}, Fabozzi Massimiliano², Reggio Stefano¹, Pannullo Mario¹, Amato Bruno³, Grimaldi Luciano¹

From 26th National Congress of the Italian Society of Geriatric Surgery
Naples, Italy. 19-22 June 2013

Abstract

Background: Rectovaginal fistulas (RVFs) have multiple causes, size and location on which the surgical treatment depends.

Description: The Authors consider different approaches to RVFs and describe a clinical case of recurrent high RVF.

Conclusions: Most RVFs can be successfully repaired, although many interventions may be necessary. A colostomy with delayed repair may improve RVFs outcome. Moreover, several authors indicate Mucosal Advancement Flap and Babcock-Bacon technique as the treatments of choice respectively for low and high RVFs (complex and recurrent) and emphasize the placement of endoscopic prosthesis in cases of difficult healing of the anastomosis.

Introduction

Rectovaginal Fistulas (RVFs) represent an arduous challenge even for the most expert surgeons can and a distressing situation for the women afflicted. Even if they only constitute about 5% of all the anorectal fistulas [1], numerous techniques have been suggested for their treatment, but unfortunately no one can assure a definitive result in the majority of cases.

The symptoms reported by majority of patients are unbearable for social, emotional, and sexual morbidity, and may sometimes be disabling. The RVF infact may cause the passage of gas and faecal material in the vagina giving out bad odor, a chronic vaginal or perineal suppuration and also the rectal syndrome with anal incontinence due to serious lesions of the anal sphincter. whose incidence in the women afflicted is very high (about 35% in the primiparous women and even 44% in the multiparous women) [2].

The anovaginal clinical examination and proctoscopy are essential for the diagnosis and definition of the anatomical and aetiological characteristics. If the RVF is of small dimension, it appears as a small depression on the vaginal mucosa, whose presence will be confirmed by a delicate probing of the passage. If the doubt persists, a tampon can be inserted in the vagina and methylene blue injected into the rectum. thus the anal orifice is closed for 15-20 min and the tampon is examined: if it is not stained, a diagnosis of RVF is very improbable.

Other investigations (for the search of collateral passages and their related complications) include: fistulography, vaginography, barium enema, pelvic C.T. with contrast medium, anal manometry and when indicated electroneuromyography for the evaluation of the pudend nerve function [3].

Clinical case

The clinical case we present concerns an incredible history of bilateral asymptomatic ovarian cysts, operated with several complications and followed by the appearance of a RVF.

R.J. is a 68 years old patient of, who is a carrier of HCV, with surgical antecedents of appendectomy (about

* Correspondence: michele.danzi@unina.it

¹Department of Specialized Surgery - Division of Gastrointestinal Surgery Rehabilitation of Election and Emergency. "Federico II" University, Naples - Italy

Full list of author information is available at the end of the article

40 years ago) and hysterectomy (about 20 years ago). Hospitalized in the department of gynecology for two tumefactions of the ovaries, in April 2007 she had a bilateral oophorectomy with some complications due to numerous and tenacious adhesive bridges, which were caused by previous surgical interventions. On this occasion, unfortunately it was not possible to avoid a lesion of the anterior wall of the Rectum, that forced the surgeon to perform the Hartman intervention. Her post-operative condition was normal. The patient was hospitalized and underwent surgery 2 months later to restore the intestinal continuity, (with the same difficulty found during the previous intervention, due to a complete freezing of the pelvis). Also on this occasion the ablation of visceral adhesions unfortunately caused bladder and small intestine lesions which were immediately sutured. During hospitalization urine leakage from the abdominal drainage tube was found, thus a re-intervention was needed. By opening the peritoneum (which was almost nonexistent after several incisions), a left ureter lesion was individuated, so a termino-terminal anastomosis on probe with double J ends (stayed there for about 4 months) was made.

The urinary problem seemed to be resolved, but 6 days later the patient presented an output of dark and maleodorous material from the vagina, associated with a fever: a diagnosis of RVF was made. The patient was sent to us by Gynecologic Department to resolve this complication.

On clinical examination the patient was in a depressed state with asthenia, "abdominal facies". By using deep palpation, we observed a plethoric and less tractable abdomen, especially in the inferior quadrants, but peristalsis was present. Other clinical parameters were: temperature 37.8°C; cardiac frequency = 72 bpm; blood pressure = 140/80 mmHg.

The clinical and instrumental examinations showed a septic state due to the presence of a pelvic abscess identified by CT, associated with rectovaginal and enterocutaneous fistulas.

We positioned a venous central catheter (v.c.c.) and made a derivative loop colostomy on the right colon. At the same time appropriate antibiotic therapy was started. After this intervention the patient's general conditions seemed to be getting better, leucocytosis was reduced, renal function returned to normal except for a chronic infection of the urinary tract. The patient was without fever and the fistula seemed to be healed, so she was discharged in August, with the necessary medical treatment. During the check-up, a month after the discharge, the clinical examination seemed to confirm the healing of fistula, but the decision of restoring digestive continuity depended on the result of the pelvic C.T. and on the barium enema. Unfortunately these

investigations confirmed the persistence of the fistula with the contrast liquid coming from the vaginal orifice, and the presence of a left pelvic subperitoneal abscess with diameter of 6 cm.

Another surgical intervention was needed to evacuate the abscess and to make a colectomy with tennino-terminal anastomosis according to Knight and Griffen. The post-operative course was normal and during the following clinical examination no vaginal or rectal side orifices were found.

Throughout the following months the patient was in good physical and psychological condition (previously numerous psychiatric consultations had been necessary to keep the patient from being depressed). The digestive function was good until the reappearance of the fistula at the fornix level, on the vaginal back wall, which is in communication with the rectum anterior wall at a distance of about 7 cm from the anus.

A surgical treatment was also needed in this occasion. After the positioning of a v.c.c., a resection of the previous colorectal anastomosis was made and a new coloanal anastomosis was performed according to Babcock-Bacon. After the mobilization of the colon, it was pulled out through the anus up to the healthy colic tract above the lesion. Then it was dissected about 3 cm above the combed line with the excision of the fistula tract and the mucosectomy of the inside sphincter. The pulled out colon was fixed by a tape to a Foley n 18 catheter and to a glass rod to prevent a retraction, which were both removed on the 3th day. The two stumps (colonic mucosa and anus derma) were anastomized on the 9th day by local anesthesia. The post-operative course was complicated by bleeding between the distal colon and retracted rectal stump, and the attempt to treat it by *electrocaugulation* failed. As anastomosis reconstruction was not possible because of the excessive stump tension due to the previous resections, we decided to position a covered self-expanding endoanal prosthesis. At the check-up 6 days later, the anal channel seemed to be healthy. The anterior part of the anastomosis was well healed over, but the posterior presented an opening of the anastomotic border with a residual bleeding due to the internal dislocation of the prosthesis toward the top. Thus a second prosthesis was placed below the first, to completely cover the anastomosis. The post-operative course was normal. Some days later the bleeding was stopped and the posterior part of the anastomosis was healing.

One year later the patient was in good general condition. The proctologic examination did not show anomalies and the prosthesis was well positioned. The Methylene Blue injection through the anus did not show any fistulas. During vaginal inspection a residual scarred depression was found on the posterior wall but no

fistulas were present. Thus confirming the healing of the wound. For these reasons we decided to close the colostomy. At the following check-up, 15 months later, the patient was free of disease.

Discussion

Recurrent RVFs are a very difficult problem for surgeons. As suggested by the clinical case described there is not only one way of treating RVFs, but the approach depends on their etiology, anatomopathologic characteristics (for example the anal sphincter involvement) and previous attempts to repair them. RVFs are classified according to etiology, location and the dimensions.

They may also be classified as simple or complex (Table 1). Various techniques have been described to repair RVFs and their associated anatomical defects [4,5]. Their classification as simple or complex fistulas, other than reflecting the different anatomical characteristics, may also indicate the best treatment to use.

The surgical techniques could be divided according to the access used:

➤ **Perineal or low access** [6]: is recommended for simple RVFs and allows techniques with:

- Direct suture

- by perineum-vaginal access according to Musset [7];

- by trans-anal access according to the Mucosal Advancement Flap (MAF) [8];

Interventions of interposition of the Inside Rectum Mescler or according to Martius's technique and its variants [9];

➤ **Combined or mixed access**: is generally recommended for complex RVFs and allows:

- direct suture with or without epiploon interposition;

- rectoplasty according to Bricker [10];

- resection of the Rectum

- with trans-anal lowering of the colon (Babcock-Bacon's Pull Through resection or according to Parks) [11,12];

- with Knight-Griffen's anastomosis;

- Miles's abdomino-perineal amputation.

Lowry [13] reported a success rate of 88% in patients undergoing a primary repair of simple RVFs by MAF. The success percentage fell to 85% with the second repair and to 55% at the third attempt. This decreased success rate may be due to the presence of inflammation, tension, hematomas or underlying diseases, thus a different approach is recommended for fistulas known to have previously failed with MAF intervention. The MAF is also suitable for patients with distal RVFs; infact excellent results have been reported with this technique by some authors [14-16].

A simple colostomy presents a low rate of spontaneous healing (about 35.3%) [17]. Traumatic RVFs could spontaneously recover after the ablation of the foreign body, the drainage of the purulent collection and the protection colostomy above the lesion. In this case it is necessary to wait at least 3-6 months before surgical re-intervention. Invasive laparotomy including re-anastomosis should never be performed and the rectal pull-through operation or abdominoperineal resection should only be reserved when other interventions fail [17]. Patients with sphincter damage should undergo sphincteroplasty using either fistulectomy or MAF as the first repair. Instead, MAF is not recommended for persistent complex fistulas. While Lowry suggests waiting for a period of 3-6 months before attempting a second repair, Hibbard affirms that waiting a period of 3 months is usually sufficient to resolve the acute state of inflammation and infection. RVFs repair of Lowry by the vaginal approach is not effective in the 90% of cases, so the trans-anal way according to Greenwald and Hoexter (Mucosal Advancement Flap) is preferred for the definitive resolution of the problem (this is obtained in 95% of cases) [8,18]. For these reasons the Greenwald's technique could represent the treatment of choice for its simplicity and for the best results. The complication rate (bleedings, infections, relapses) is very low (15%) in relationship to the other techniques proposed (Table 2). For high and complex RVFs the treatment of choice has not been fixed yet due to poor documentation is present in literature. Generally, the abdominal or abdominoperineal approach is preferable for these type

Table 1 RVFs Classification [24]

AETIOLOGY:

- Congenital

- Acquired

- Traumatic (obstetric, post-operative, caused by rape or foreign bodies)

- Infectious (perirectal abscesses, diverticulitis, tuberculosis, lymphogranuloma venereum, Bartholin gland abscesses)

- Chronic inflammatory Bowel Diseases (Crohn's Disease and Ulcerative Colitis) Post-radiotherapy

- Neoplasms (primary, recurrent, metastatic)

- Idiopathic

LOCATION:

- High (With the orifice fistulous to the level of back fornix)

- Mid (between uterine cervix and the vaginal fork)

- Low (to the level of the vaginal fork): divided into oversphinteric and intrasphinteric

SIMPLE: Low or with diameter < 2.5 cm (traumatic or infectious)

COMPLEX: High with diameter > 2.5 cm (Inflammatory Bowel Diseases, post-radiotherapy, neoplasms).

of fistulas (neoplastic, atinic or in course of inflammatory bowel diseases RVFs) [19,20]. RVFs caused by inflammatory bowel diseases (IBD), have not possibility of spontaneous recovery, even if they are treated with antimetabolites therapies (Cyclosporine, 6-Mercaptopurine + Metronidazole, etc.) [21]. The same course have atinic or neoplastic fistulas. Simple RVFs have surely a better prognosis than high RVFs for which complex and aggressive techniques are necessary. A pre-operative evaluation of the fistula's tract and its careful excision is very important for a successful treatment. It is also necessary to take multiple biopsies of the borders to exclude other possible underlying diseases. A functional evaluation of the anal sphincter using transanal ultrasonography, manometry and electromyography is also needed, in fact it may be damaged by the fistulous tract or during surgical interventions. In the case described techniques as manometry and endoscopy showed a normal sphincteric function and the absence of other lesions either in the pre- or in post-operative time. The MAF technique has not good results in complex or recidivant fistulas [22]. For these types of fistulas, the

pull-through resection with coloanal anastomosis on 7-9th day assures a complete excision of the fistulous tract. This operation, associated to an endoscopic positioning of the prothesis and a total parenteral nutrition for 14 days, allows the recovery of the general patient's condition of the patient. In some cases, a preventive intestinal derivation for at least 3-6 months is necessary and is the only possible and definitive therapy.

Conclusions

This clinical case suggests us some important conclusions:

1. High RVFs therapy is not unique, but depends on different factors;
2. Surgery assure the complete excision of the fistulous tract;
3. The importance of a colostomy for the anastomosis protection, especially after reefing approach when the rectal mucosa is already damaged [23].

In our case the opening on the anastomosis and the fibrous retraction of the stumps with bleeding were treated infact with lowering the colic stump wasn't possible for its difficult isolation. We used an *autoexpandible covered endoprothesis* for the semplicity of positioning and for the therapeutic effect (hemostasis and prophylaxis of recidivant fistula); in fact, for a successful surgical repair, it is necessary to respect some principles such as handling and gentle dissection of tissues, a correct debridement and the colon mobilization to make tension-free anastomosis. Unfortunately no one is possible in our case.

We also want to confirm the MAF as the treatment of choice for the simple fistulas and not for the recidivant. For complex and recidivant fistulas, according to our experience, the methodic of the pull-through resection may have good results for the possibility of a radical excision. The reason for this aggressive approach is the lower rate of healing after a primary unsuccessful repair.

The endoscopic approach besides, in this specific case, was providential, infact it provided the protection of the rectal mucosa and in the same time the stop of the anastomotic collar bleeding. Thus we may propose the positioning of self-expandible covered endoprothesis when a protection of the rectal mucosa is needed. Simple RVFs repair by MAF has excellent results, while complex RVFs have to must be treated by more aggressive methods. In fact, once the first attempt to repair by derivative colostomy has failed, a difficult intervention such as the pull-through resection ace. to Babcock-Bacon must be managed by expert surgeons. Although in most cases repeated surgery may be necessary, the pull through resection is sure, radical and effective whereas other techniques have failed.

Table 2 INDICATIONS OF SURGICAL TECHNIQUES

AGE:

- Young women: less aggressive procedures with safeguard of vaginal integrity, when it is possible
- Elderly women: interventions for the drastic resolution of the problem with abdominoperineal approach;

ETIOLOGY:

- Post operative, traumatic and infectious RVFs
 - MAF;
 - recidivists: bowel resection with trans-anal anastomosis acc.to Babcock (or to Parks) + epiploon interposition;
- Post-radiotherapy RVFs
 - colostomy + waiting for 3-6 months: if it's necessary, bowel resection with coloanal anastomosis or pull-through resection acc. To Babcock-Bacon;
 - in absence of bleeding: bulbocavernosus muscle interposition am to Martius;
 - in case of vulva sclerosis: inside rectum muscle interposition;
 - when other attempt to repair fail -> definitive colostomy.
- IBD RVFs
 - drainage of perinal abscesses o trans-sphincteric loop;
 - if it's necessary: colostomy for 3-6 months;
 - 3 months of antimetabolites therapy;
 - in more serious cases: bowel resections.
- Tumoral RVFs
 - low anterior resection of the rectum or amputation of Miles + back colectomy;

LOCALIZATION:

- High RVFs requires interventions with combined approach;
- Low RVFs requires a perineal approach
- Mid RVFs require an approach depending on etiology and eventually associated lesions

List of abbreviations

RVF: recto-vaginal fistula; CT: computed tomography; HCV: hepatitis C virus; vcc: venous central catheter; MAF: mucosal advancement flap; IBD: inflammatory bowel disease.

Competing interests

The authors declare that they have no competing interests.

Authors' contributions

D.M.: conception and design, interpretation of data, given final approval of the version to be published.

F.M.: acquisition of data, drafting the manuscript, given final approval of the version to be published.

R.S.: acquisition of data, drafting the manuscript, given final approval of the version to be published.

P.M.: acquisition of data, drafting the manuscript, given final approval of the version to be published.

A.B.: critical revision, interpretation of data, given final approval of the version to be published

G.L.: conception and design, critical revision, given final approval of the version to be published.

Authors' information

DM: Assistant Professor of Surgery at University Federico II of Naples.

FM: MD, PhD, in Surgery at U. Parini Hospital of Aosta.

RS: Resident in Surgery at University Federico II of Naples.

PM: Resident in Surgery at University Federico II of Naples.

AB: Associate Professor of Surgery at University "Federico II" of Naples.

GL: MD, PhD, in Surgery at University Federico II of Naples.

Declarations

Funding for this supplement has come from personal funds.

This article has been published as part of *BMC Surgery* Volume 13 Supplement 2, 2013: Proceedings from the 26th National Congress of the Italian Society of Geriatric Surgery. The full contents of the supplement are available online at <http://www.biomedcentral.com/bmc/surg/supplements/13/S2>

Authors' details

¹Department of Specialized Surgery - Division of Gastrointestinal Surgery Rehabilitation of Election and Emergency. "Federico II" University, Naples - Italy.

²Department of General Surgery. "U. Parini" Hospital, Aosta - Italy.

³Department of General, Geriatric, Oncologic Surgery and Advanced Technologies. "Federico II" University, Naples - Italy.

Published: 8 October 2013

References

1. Khanduja KS, Padmanabhan A, Kerner B, *et al*: **Reconstruction of rectovaginal fistula with sphincter disniption by combining rectal mucosal advancement flap and anal sphincteroplasty.** *Dis Colon Rectum* 1999, **42**(11):1432-1437.
2. Milito G, Pisani A, Venditti D, *et al*: **The endorectal repair of rectovaginal fistulae.** *Minerva Chir* 1999, **54**(3):191-194.
3. Mazier WP, Senagore AJ, Schiesel EC: **Operative repair of anovaginal and rectovaginal fistulas.** *Dis Colon Rectum* 1995, **38**:4-6.
4. Tozer PJ, Balmforth D, Kayani B, Rahbour G, Hart AL, Phillips RK: **Surgical management of rectovaginal fistula in a tertiary referral centre: many techniques are needed.** *Colorectal Dis* 2013, Jan 19.
5. Schlöricke E, Zimmermann M, Hoffmann M, Laubert T, Nolde J, Hildebrand P, Bruch HP, Bouchard R: **Surgical treatment and prognosis of rectovaginal fistulae according to their origin.** *Zentralbl Chir* 2012, **137**(4):390-395, Aug.
6. Schloericke E, Hoffmann M, Zimmermann M, Kraus M, Bouchard R, Roblick UJ, Hildebrand P, Nolde J, Bruch HP, Limmer S: **Transperineal omentum flap for the anatomic reconstruction of the rectovaginal space in the therapy of rectovaginal fistulas.** *Colorectal Dis* 2012, **14**(5):604-610, May.
7. Musset R: **Fistules recto-vaginales.** *Encycl Med Chir Techn Chir. Urol-Gynéc* 41-870, 4.4.06: 6p.
8. Greenwald JC, Hoexter B: **Repair of rectovaginal fistula.** *Surg Gynecol Obstet* 1978, **146**:443-445.
9. Boronow RC: **Repair of the radiation-induced vaginal fistula utilizing the Martins technique.** *World J Surg* 1986, **10**:237-48.
10. Bricker EM, Johnston WD, Patwardhan RV: **Repair of postirradiation damage to colorectum: A progress report.** *Ann Surg* 1981, **193**:555-564.
11. Parks AG, Allen CL, Frank JD, *et al*: **A method of treting post-irradiation rectovaginal fistulas.** *Br J Surg* 1978, **65**:417-421.
12. Cutait R, Ioshimoto M, *et al*: **Abdominoperineal endoanal pull-through resection.** *Dis Colon Rectum* 1985, **28**:294-299.
13. Lowry AC, Thorson AG, Rothenberger DA, *et al*: **Repair of simple rectovaginal fistulas: influence of previous repairs.** *Dis Colon Rectum* 1988, **31**:676-678.
14. De Parades V, Dahmani Z, Blanchard P, Zeitoun JD, Sultan S, Atienza P: **Endorectal advancement flap with muscular plication: a modified technique for rectovaginal fistula repair.** *Colorectal Dis* 2011, **13**(8):921-925, Aug.
15. Kodner Mazor A, Shemesh El, *et al*: **Endorectal advancement flap repair of rectovaginal and other complicated anorectal fistulas.** *Surgery* 1993, **114**:682-690.
16. Wise WE Jr, Aguilar PS, Padmanabhan A, *et al*: **Surgical treatment of low rectovaginal fistulas.** *Dis Colon Rectum* 1991, **34**:271-4.
17. Rex JC Jr, Khubchandani IT: **Rectovaginal fistula: complication of low anterior resection.** *Dis Colon Rectum* 1992, **35**:354-356.
18. Hoexter B, Labow SB, Moseson MD: **Transanal rectovaginal fistula repair.** *Dis Colon Rectum* 1985, **28**:572-575.
19. Sonoda T, Hull T, Piedmonte MR, Fazio VW: **Outcomes of primary repair of anorectal and rectovaginal fistulas using the endorectal advancement flap.** *Dis Colon Rectum* 2002, **45**(12):1622-1628.
20. Lindsey I, Smilgin-Humphreys MM, Cunningham C, *et al*: **A randomized, controlled trial of fibrin glue vs conventional treatment for anal fistula.** *Dis Colon Rectum* 2002, **45**(12):1608-1615.
21. De la Poza G, López-Sanroman A, Taxonera C, Marín-Jimenez I, Gisbert JP, Bermejo F, Opio V, Muriel A: **Genital fistulas in female Crohn's disease patients: clinical characteristics and response to therapy.** *J Crohns Colitis* 2012, **6**(3):276-280, Apr.
22. Halverson LA, Tracy LH, Fazio VW, *et al*: **Repair of recurrent rectovaginal fistulas.** *Surgery* 2001, **130**:753-758.
23. Saclarides TJ: **Rectovaginal fistula.** *Surg Clin North Am* 2002, **8**(6):1261-1272, Dec.
24. Frileux P, Berger A, Zinzindohoue F, *et al*: **Fistules rectovaginales de l'adulte.** *Ann Chir* 1994, **48**(5):412-420.

doi:10.1186/1471-2482-13-S2-S7

Cite this article as: Danzi *et al*: Surgical mistake causing an high rectovaginal fistula. A case report with combined surgical and endoscopic approach: therapeutic considerations. *BMC Surgery* 2013 **13**(Suppl 2):S7.

Submit your next manuscript to BioMed Central and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit

